



Product Name: CONFIDOR 200 SC INSECTICIDE
APVMA Approval No: 50548/124741

Label Name:	CONFIDOR 200 SC INSECTICIDE
-------------	-----------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	200 g/L IMIDACLOPRID
-------------------------	----------------------

Mode of Action:	GROUP 4A INSECTICIDE
-----------------	----------------------

Statement of Claims:	1 L For the control of various insect pests of fruit, vegetables and ornamentals as specified in the DIRECTIONS FOR USE table 10, 20 and 60 L For the control of various insect pests of cotton, fruit, vegetables and ornamentals as specified in the DIRECTIONS FOR USE table 50 L For the control of various insect pests of Brassica vegetables when applied as a soil treatment as specified in the DIRECTIONS FOR USE table
----------------------	--

Net Contents:	10L 20L 50L 60L 1L
---------------	--------------------------------

Restraints:	
-------------	--

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	<p>WITHHOLDING PERIODS</p> <p>1 L</p> <p>Harvest:</p> <p>Apples: NOT REQUIRED WHEN USED AS DIRECTED</p> <p>Cucurbits: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION</p> <p>Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION</p> <p>Brassiccas, capsicum, eggplant, potatoes, sweet potatoes: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION</p> <p>Stone fruit: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION</p> <p>Grazing:</p> <p>DO NOT graze any treated area, or cut for stock food.</p> <p>DO NOT feed produce harvested from treated area to animals, including poultry.</p> <p>10, 20, 60 L</p> <p>Foliar spray applications</p> <p>Harvest:</p> <p>Cucurbits: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION</p> <p>Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION</p> <p>Brassiccas, capsicum, eggplant, potatoes, sweet potatoes: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION</p> <p>Stone fruit: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION</p> <p>Cotton: DO NOT HARVEST FOR 13 WEEKS AFTER APPLICATION</p> <p>Soil or seedling drench applications</p> <p>Harvest:</p> <p>Apples: NOT REQUIRED WHEN USED AS DIRECTED</p> <p>Brassiccas: NOT REQUIRED WHEN USED AS DIRECTED</p> <p>Chicory, endive, lettuce (excluding head lettuce) and radicchio: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION</p> <p>Head lettuce: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION</p> <p>All applications</p> <p>Grazing:</p> <p>DO NOT GRAZE ANY TREATED AREA, OR CUT FOR STOCK FOOD.</p> <p>DO NOT FEED PRODUCE HARVESTED FROM TREATED AREA TO ANIMALS, INCLUDING POULTRY.</p> <p>50 L</p> <p>Harvest:</p> <p>Brassiccas: NOT REQUIRED WHEN USED AS DIRECTED</p> <p>Grazing:</p> <p>DO NOT graze any treated area, or cut for stock food.</p> <p>DO NOT feed produce harvested from treated area to animals, including poultry.</p>
----------------------	--

Trade Advice:	<p>EXPORT OF TREATED PRODUCE</p> <p>Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Confidor. If you are growing edible produce for export, please check with Bayer Crop Science for the latest information on MRLs and import tolerances before using Confidor.</p>
---------------	--

General Instructions:	This section contains file attachment.
-----------------------	--

Resistance Warning:	<p>INSECTICIDE RESISTANCE WARNING GROUP 4A INSECTICIDE</p> <p>For insecticide resistance management, Confidor is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Confidor and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Confidor and other Group 4A insecticides are used repeatedly. The effectiveness of Confidor on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of Confidor to control resistant insects. Confidor may be subject to specific resistance management strategies. For further information contact your local supplier, Bayer Crop Science representative or local agricultural department agronomist.</p> <p>Resistance Management Strategies</p> <p>1 L</p> <p>Aphids, whitefly and melon thrips in various crops (foliar sprays)</p> <p>Do not apply Confidor (or other group 4A insecticides) in consecutive sprays within and between seasons. Rotate with registered insecticides from other mode of action groups. Confined environments such as glasshouses:</p> <p>Annuals - Do not apply more than one spray of Confidor (or other group 4A insecticides) to any one crop.</p> <p>Perennials - Rotate with registered insecticides from other groups. Use a maximum of three Confidor (or other group 4A insecticides) sprays in any 12 month period.</p> <p>10, 20, 60 L</p> <p>Cotton aphid in cotton</p> <p>Observe the cotton industry Insecticide Resistance Management Strategy (IRMS).</p> <p>Aphids, whitefly and melon thrips in various crops (foliar sprays)</p> <p>Do not apply Confidor (or other group 4A insecticides) in consecutive sprays within and between seasons. Rotate with registered insecticides from other mode of action groups. Confined environments such as glasshouses:</p> <p>Annuals - Do not apply more than one spray of Confidor (or other group 4A insecticides) to any one crop.</p> <p>Perennials - Rotate with registered insecticides from other groups. Use a maximum of three Confidor (or other group 4A insecticides) sprays in any 12 month period.</p> <p>Brassica and lettuce (seedling drench applications)</p> <p>DO NOT apply Confidor 200 SC or any other Group 4A Insecticide following a seedling drench application of Confidor 200 SC or any Group 4A Insecticide in that crop.</p> <p>50 L</p> <p>DO NOT apply Confidor 200 SC or any other Group 4A Insecticide following a furrow spray or plant hole drench application of Confidor 200 SC or any Group 4A Insecticide in that crop.</p>
---------------------	---

Precautions:	<p>PRECAUTIONS</p> <p>10, 20, 60 L</p> <p>Seedling drench application:</p> <p>At the nursery - gloves should be worn when handling treated seedling trays or seedlings. Cotton overalls buttoned to the neck and wrist (or equivalent clothing) should also be worn, particularly if seedlings and/or trays are still wet following treatment.</p> <p>On farm including during planting - gloves should be worn when handling treated seedling trays or seedlings.</p> <p>50 L</p> <p>Furrow spray application: At planting, steps should be taken to ensure workers do not contact treated soil.</p>
--------------	--

Plant hole drench application: At planting and when applying drench, steps should be taken to ensure workers do not contact treated soil or drench mixture.

Protections:	<p>PROTECTION OF LIVESTOCK Dangerous to bees. DO NOT spray any plants in flower if bees may forage on treated flowers. DO NOT graze any treated area, or cut for stock food. DO NOT feed produce harvested from treated area to animals, including poultry.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT 1 L Imidacloprid is toxic to certain aquatic species. DO NOT contaminate streams, rivers, wetlands or watercourses with this product or used containers. Aerial application: A spraydrift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practices Manual.</p> <p>10, 20, 60 L Imidacloprid is toxic to certain aquatic species. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Aerial application: A spraydrift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practices Manual. Seedling drench application: DO NOT allow run-off from treatment areas to enter drains and waterways.</p> <p>50 L Imidacloprid is toxic to certain aquatic species. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Furrow spray and plant hole drench application (brassicas): Application should be planned to avoid run-off within 48 hours of application. Application should not be made to wet/waterlogged soils. Application is not to be performed if heavy rains are expected to occur within 48 hours. Irrigation within 48 hours of application should be minimal, and sufficient to reduce seedling stress only. Irrigation run-off from treated areas should be prevented from entering drains and waterways.</p>
--------------	--

Storage and Disposal:	<p>STORAGE AND DISPOSAL Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. 1, 10, 20, 50, 60 L non-returnable Triple rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. Do not re-use empty container for any other purpose.</p> <p>50, 60 L returnable only If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products.</p>
-----------------------	---

Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of Bayer CropScience Pty Ltd.

Safety Directions:	SAFETY DIRECTIONS Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length chemical resistant gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.
--------------------	---

First Aid Instructions:	FIRST AID If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26).
-------------------------	--

First Aid Warnings:	
---------------------	--

1 L
DIRECTIONS FOR USE

FOLIAR SPRAY APPLICATIONS

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Stone fruit	Green peach aphid, black peach aphid	Dilute spraying 25 mL/100 L Concentrate spraying Refer to the Application section	21 days	Apply at first sign of aphid infestation. Apply as a full cover spray, ensuring thorough coverage. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Do not use in equipment that requires rates greater than 125 mL/100 L of water (i.e. greater than 5 X concentrate).
Cucurbits	Green peach aphid	25 mL/100 L or 300 mL/ha	1 day	Apply at first sign of aphid infestation.
Capsicum, eggplant, potato			7 days	
Tomato			3 days	
Brassicas	Grey cabbage aphid, turnip aphid	7 days	Apply at first sign of aphid infestation. Add a wetting agent.	
Sweet potato	Silverleaf whitefly, including type B	25 mL/100 L or 250 mL/ha	7 days	Apply at first sign of whitefly or melon thrips infestation. Apply dilute sprays (25 mL/100 L) to run off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Cucumber			1 day	
Egg plant			7 days	
Roses	Aphids	25 mL/100 L	—	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental plants	Aphids, azalea lace bug, bronze orange bug, citrus mealybug, Fuller's rose weevil, greenhouse thrips, harlequin bug, silverleaf whitefly (suppression)			
	Hibiscus flower beetle			Spray buds and flowers as needed.
	Longtailed mealybug			Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids			Spray at first sign and then a week later.
	Soft scales			Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green peach aphid	25 mL/100 L		Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.

Confidor 200 SC Insecticide

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Pandanus trees	Flatid (<i>Jamella australiae</i>)	Spot spray 875 mL/100 L of water Stem injection 1.75 L/1 L of water		Spot spray: Spray 100 mL of mixture directly into the leafy throat of each head. Stem injection: Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Confidor, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

SOIL APPLICATIONS

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Apples	Woolly aphid	Chemical control 6 mL/1 L of water/tree Beneficial insect plus chemical control (e.g. <i>Aphelinus mali</i> plus Confidor) 3 mL/1 L of water/tree	-	For trees up to 7 years of age. During late summer or autumn, apple trees with woolly aphid colonies or damage should be identified and marked for treatment the following season. At green tip to petal fall, apply 1 litre of the prepared Confidor mixture to moist soil immediately around the base of the tree trunk. Ensure the mixture infiltrates the soil around the trunk and does not run off the soil. Control weeds before application. Do not disturb or remove the soil around the trunk during the season. If aerial colonies are present at application, maximum effectiveness may not be achieved until the following season. Do not treat more than once in any 2 year period.
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot		Use as a soil drench for pots up to 20 L capacity. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Confidor mixture, and then water it in well immediately after application.
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height		Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400-500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7-10 days after treatment. Treat at least 6-10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water		Use as a soil drench. 5 L of mixture will treat twenty 6 L pots. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Confidor mixture, and then water it in well immediately after application.
Roses	Aphids	3.5 mL/2 L water/plant		Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Confidor to the 2 L of water. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Confidor mixture, and then water it in well immediately after application.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

10, 20, 60 L
DIRECTIONS FOR USE

FOLIAR SPRAY APPLICATIONS

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Cotton	Aphids	250 mL/ha + Pulse Penetrant at 0.2% v/v (2 mL/L water) or equivalent organosilicone surfactant* (e.g. Freeway Gold) * Check with Bayer Crop Science for recommended organosilicone surfactants	13 weeks	<p>The addition of Pulse Penetrant or equivalent is critical for the performance of Confidor. Apply early in the establishment of an aphid infestation when numbers are low (i.e. no more than 1 or 2 leaves per plant with honeydew present). Applications made later than this may result in reduced control.</p> <p>Shorter residual control may be evident and a repeat application of a registered aphicide (follow the Cotton Insecticide Resistance Management Strategy for cotton aphid) may be required to achieve complete control:</p> <ul style="list-style-type: none"> • if applications of Confidor plus Pulse or equivalent are timed too late (see above); or • if existing high density aphid colonies (hotspots) are present; or • if aphids have established throughout the plant canopy (especially lower in the canopy); or • if there is high reinestation pressure; or • if there is rapid crop growth; or • if Confidor plus Pulse or equivalent is used following a spray-failure (e.g. resistance to organophosphate or carbamate insecticides). <p>Note: Where resistance to carbamates or organophosphates is suspected, Confidor plus Pulse or equivalent should be used first so as not to delay control of the aphids present.</p>
	Mirids, brown flea beetle			Apply when pest numbers reach treatment threshold levels as determined by field checks.
Stone fruit	Green peach aphid, black peach aphid	Dilute spraying 25 mL/100 L Concentrate spraying Refer to the Application section	21 days	Apply at first sign of aphid infestation. Apply as a full cover spray, ensuring thorough coverage. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Do not use in equipment that requires rates greater than 125 mL/100 L of water (i.e. greater than 5 X concentrate).
Cucurbits	Green peach aphid	25 mL/100 L or 300 mL/ha	1 day	Apply at first sign of aphid infestation.
Capsicum, eggplant, potato			7 days	
Tomato			3 days	
Brassiccas	Grey cabbage aphid, turnip aphid	7 days	Apply at first sign of aphid infestation. Add a wetting agent.	

Confidor 200 SC Insecticide

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Sweet potato	Silverleaf whitefly, including type B	25 mL/100 L or 250 mL/ha	7 days	Apply at first sign of whitefly or melon thrips infestation. Apply dilute sprays (25 mL/100 L) to run off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Cucumber			1 day	
Egg plant	Melon thrips		7 days	
Roses	Aphids	25 mL/100 L	—	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental plants	Aphids, azalea lace bug, bronze orange bug, citrus mealybug, Fuller's rose weevil, greenhouse thrips, harlequin bug, silverleaf whitefly (suppression)			
	Hibiscus flower beetle			Spray buds and flowers as needed.
	Longtailed mealybug			Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids			Spray at first sign and then a week later.
	Soft scales	25 mL/100 L		Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green peach aphid			Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Pandanus trees	Flatid (<i>Jamella australiae</i>)	Spot spray 875 mL/100 L of water Stem injection 1.75 L/1 L of water		Spot spray: Spray 100 mL of mixture directly into the leafy throat of each head. Stem injection: Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Confidor, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

Confidor 200 SC Insecticide

SOIL APPLICATIONS

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Apples	Woolly aphid	Chemical control 6 mL/1 L of water/tree Beneficial insect plus chemical control (e.g. <i>Aphelinus mali</i> plus Confidor) 3 mL/1 L of water/tree	-	For trees up to 7 years of age. During late summer or autumn, apple trees with woolly aphid colonies or damage should be identified and marked for treatment the following season. At green tip to petal fall, apply 1 litre of the prepared Confidor mixture to moist soil immediately around the base of the tree trunk. Ensure the mixture infiltrates the soil around the trunk and does not run off the soil. Control weeds before application. Do not disturb or remove the soil around the trunk during the season. If aerial colonies are present at application, maximum effectiveness may not be achieved until the following season. Do not treat more than once in any 2 year period.
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot		Use as a soil drench for pots up to 20 L capacity. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Confidor mixture, and then water it in well immediately after application.
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height		Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400-500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7-10 days after treatment. Treat at least 6-10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water		Use as a soil drench. 5 L of mixture will treat twenty 6 L pots. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Confidor mixture, and then water it in well immediately after application.
Roses	Aphids	3.5 mL/2 L water/plant		Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Confidor to the 2 L of water. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Confidor mixture, and then water it in well immediately after application.

SEEDLING DRENCH APPLICATIONS**RESTRAINT:**

DO NOT transplant seedlings treated by seedling drench into hydroponic production systems.

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Brassicas (broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi)	Silverleaf whitefly, including type B	Seedling drench 40 mL/1000 seedlings	-	BRASSICAS ONLY Green peach aphid, onion thrips: When Confidor is used for the control of silverleaf whitefly, including type B, control of green peach aphid and onion thrips will also be achieved.
	Green peach aphid, onion thrips	Seedling drench 20-40 mL/1000 seedlings		
Lettuce (head lettuce)	Currant-lettuce aphid	Planting densities up to 70,000 plants/ha 35-55 mL/1000 seedlings applied as a seedling drench	6 weeks	Apply to the transplant cell, once only, prior to transplanting.
Lettuce (excluding head lettuce), chicory, endive, radicchio		Planting densities from 70,000 to 110,000 plants/ha 12.5-35 mL/1000 seedlings applied as a seedling drench	4 weeks	For applications specifically against green peach aphid and onion thrips, select the rate taking into account the likely crop type, crop growing period, the anticipated degree of pest infestation and previous field experience (e.g. consideration of crop variety, time of year, predator activity, soil type). Refer to <u>Additional Critical Comments – all crops</u> below for further advice.
		Planting densities from 110,000 to 150,000 plants/ha 12.5-25 mL/1000 seedlings applied as a seedling drench		LETTUCE, CHICORY, ENDIVE, RADICCHIO ONLY Seedling drench (prior to transplanting) Apply to the transplant cell, once only, prior to transplanting. The maximum recommended application rate per 1000 seedlings will depend on the proposed field planting density, as shown in the RATE column.
		Planting densities up to 70,000 plants per hectare 12.5 mL/1000 seedlings applied as a seedling drench		Planting densities up to 70,000 plants per hectare In the RATE column, identify the desired planting density and the corresponding recommended rate range. Select the rate taking into account the likely crop growing period, the anticipated degree of pest infestation and previous field experience (e.g. consideration of lettuce type, variety, time of year, predator activity, soil type).
				The lower rate will usually be adequate for crops with a short growing period (e.g. less than 6-7 weeks), while the higher rate will be necessary for most situations where conditions are highly favourable for aphid infestation and for longer crop growing periods (e.g. greater than 6-7 weeks). The lower rate may also be appropriate where control from the Confidor application is not required for the entire crop growing period and where insecticides with alternative modes of action will be used as part of a broader lettuce aphid management strategy

Confidor 200 SC Insecticide

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
				<p>during latter stages of crop development.</p> <p>Refer to <u>Additional Critical Comments – all crops</u> below for further advice.</p> <p><u>Planting densities from 70,000 to 110,000 and 110,000 to 150,000 plants per hectare</u></p> <p>In the RATE column, identify the desired planting density and the corresponding recommended rate range.</p> <p>Select the rate taking into account the likely crop growing period, the anticipated degree of pest infestation and previous field experience (e.g. consideration of lettuce type, variety, time of year, predator activity, soil type).</p> <p>The lower rate will be suitable where a short duration of control is required, while a higher rate will generally provide an increased length of control.</p> <p>However these rates may not provide control of currant lettuce aphid for the entire crop growing period. Monitor crops and if required use insecticides with alternative modes of action as part of a broader lettuce aphid management strategy during latter stages of crop development.</p> <p>Refer to <u>Additional Critical Comments – all crops</u> below for further advice.</p> <p><u>Planting densities from 150,000 up to 300,000 plants per hectare</u></p> <p>In the RATE column, identify the desired planting density and the corresponding recommended rate.</p> <p>This rate may not provide control of currant lettuce aphid for the entire crop growing period. Monitor crops and if required use insecticides with alternative modes of action as part of a broader lettuce aphid management strategy during latter stages of crop development.</p> <p>Refer to <u>Additional Critical Comments – all crops</u> below for further advice.</p> <p><u>Additional Critical Comments – all crops</u></p> <p>Confidor provides effective management of pest populations. However, Confidor may not provide complete control of pests for the entire growing period in all situations. Crops should be monitored for pests following transplanting and throughout the life of the crop. If pests are observed in the crop additional chemical control may be required, in which case an insecticide with a different mode of action should be used. Refer to GENERAL INSTRUCTIONS for Resistance Management</p>

Confidor 200 SC Insecticide

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
				<p>Strategy information.</p> <p>Refer to GENERAL INSTRUCTIONS for Seedling Drench Application directions and PRECAUTIONS when handling treated seedlings.</p> <p>Seedling damage may result from Confidor seedling drench treatment particularly if transplanting does not occur soon after treatment. It is recommended that transplanting occur within 24 hours of treatment and that planted seedlings receive sufficient irrigation (preferably using overhead sprinklers) as soon as possible after transplanting to further minimise the risk of seedling damage. This may be particularly relevant under conditions of rapid drying of the transplant cell medium.</p> <p>If watering is required between applications and planting, it should be done sparingly, only as required. DO NOT allow run-through from the cells.</p> <p>When transplanting treated seedlings ensure that the growing medium is fully transferred to the field with each seedling.</p> <p>Seedling production nurseries supplying Confidor treated seedlings must ensure that;</p> <ul style="list-style-type: none"> a) Supplied batches of seedlings are clearly identified as having been treated with Confidor 200 SC Insecticide. b) Paperwork accompanying the seedlings and provided to the recipient indicates the rate of Confidor applied per 1000 seedlings, and the time and date of treatment. c) Growers accepting delivery of treated seedlings have obtained a copy of <i>Confidor 200 SC seedling drench – instructions for growers</i>, available from Bayer Crop Science (1800 804 479 or www.crop.bayer.com.au)

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

50 L**DIRECTIONS FOR USE**

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Brassicas (broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi)	Silverleaf whitefly, including type B	Plant hole drench, furrow spray 60 mL/1000 seedlings	-	<p>Green peach aphid, onion thrips: When Confidor is used for the control of Silverleaf whitefly, including type B, control of green peach aphid and onion thrips will also be achieved.</p> <p>Confidor provides effective management of pest populations. However, Confidor may not provide complete control of pests for the entire growing period. Crops should be monitored for pests following planting and throughout the life of the crop. If pests are observed in the crop additional chemical control may be required, in which case an insecticide with a different mode of action should be used. Refer to GENERAL INSTRUCTIONS for resistance management strategy information.</p> <p>Plant hole drench (at or post-transplanting) Mix Confidor with sufficient water to allow a constant volume of at least 50 mL of drench mixture per plant. Apply the selected volume of drench mixture in the planting hole at planting or within 2 days after planting.</p> <p>Furrow spray (prior to planting) Mix Confidor with water, using at least 2 litres of spray mixture per 100 m of row. Apply to open furrow not earlier than 5 days prior to planting as a narrow band of spray centred under the plant row. Do not leave the sprayed band exposed to sunlight; sprayed soil should be covered immediately. After final shaping of the planting bed, the treated layer of soil should be approximately 100 mm below the soil surface. This method of application may provide less effective control than seedling drench or plant hole drench application methods.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

1 L GENERAL INSTRUCTIONS

Mixing

Prior to pouring, shake container vigorously, then add the required quantity of Confidor 200 SC to water in the spray vat while stirring or with agitators in motion.

Application

Foliar Spray Application

Where rates are expressed in mL/100 L mix specified rates of product per 100 L water and spray plants to "point of run-off" stage, thoroughly covering all plant surfaces.

Where rates are expressed in mL/ha apply specified rates per hectare using appropriate spray equipment and sufficient water to provide adequate penetration and coverage. Aim to apply even small droplet coverage over all plant surfaces. Equipment settings and water volume may need to vary, depending on the growth stage of the crop.

Special Instructions for Stone Fruit

Dilute Spraying

- ◆ Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- ◆ The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- ◆ Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- ◆ The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- ◆ Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- ◆ Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- ◆ The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500 L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is: 3 X (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$)
4. As the dilute label rate is 25 mL/100 L for stone fruit, then the concentrate rate becomes 3×25 , that is 75 mL/100 L of concentrate spray.

- ◆ The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- ◆ Do not use a concentrate rate higher than that specified in the Critical Comments.
- ◆ For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Note on Ornamentals

Confidor 200 SC Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

COMPATIBILITY

Confidor 200 SC is compatible with Antracol® and Baycor®. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Bayer Crop Science, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Bayer Crop Science cannot be responsible for the behaviour of such mixtures.

10, 20, 60 L

Mixing

Prior to pouring, shake container vigorously, then add the required quantity of Confidor 200 SC to water in the spray vat while stirring or with agitators in motion.

Application

Foliar Spray Application

Where rates are expressed in mL/100 L mix specified rates of product per 100 L water and spray plants to "point of run-off" stage, thoroughly covering all plant surfaces.

Where rates are expressed in mL/ha apply specified rates per hectare using appropriate spray equipment and sufficient water to provide adequate penetration and coverage. Aim to apply even small droplet coverage over all plant surfaces. Equipment settings and water volume may need to vary, depending on the growth stage of the crop.

Special Instructions for Cotton

Thorough coverage of cotton plants is essential to achieve maximum performance from Confidor plus Pulse. Equipment should be calibrated to achieve a **minimum of 60 droplets/cm² on the target foliage**. A droplet **Volume Median Diameter (VMD)** for optimum performance from Confidor plus Pulse is dependent on equipment and is defined below. Do not apply when unfavourable environmental conditions may reduce the quality of spray coverage.

Ground Application (Cotton)

Application using ground equipment should be made using **hollow cone nozzles** with a **minimum spray volume of 100 L/ha**. Hollow cone nozzles are recommended but if flat fan nozzles are used, higher water volumes will be required and nozzles should be configured to ensure thorough coverage. A droplet **VMD of 150 - 180 microns** must be used. Where multiple nozzles per row are used, they should be of the same specification to ensure that each nozzle contributes an equal proportion of the required dose. Where multiple nozzles per row are used (particularly for banded applications) ensure the correct nozzle overlap pattern is achieved on the target foliage. **Banded applications less than 100% are not recommended beyond the 15 node crop stage.**

Aerial Application (Cotton)

Apply in a **minimum spray volume of 25 L/ha**. A droplet **VMD of 120 - 150 microns** must be used. Do not exaggerate swath width or exceed a swath width of 20 to 22 m. Do not apply Confidor plus Pulse using Ultra Low Volume (ULV) methods. The use of large droplet placement equipment is not recommended.

Special Instructions for Stone Fruit

Dilute Spraying

- ◆ Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- ◆ The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- ◆ Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.

- ◆ The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- ◆ Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- ◆ Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- ◆ The mixing rate for concentrate spraying can then be calculated in the following way:
EXAMPLE ONLY
 1. Dilute spray volume as determined above: For example 1500 L/ha
 2. Your chosen concentrate spray volume: For example 500 L/ha
 3. The concentration factor in this example is: 3 X (i.e. 1500 L ÷ 500 L = 3)
 4. As the dilute label rate is 25 mL/100 L for stone fruit, then the concentrate rate becomes 3 x 25, that is 75 mL/100 L of concentrate spray.
- ◆ The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- ◆ Do not use a concentrate rate higher than that specified in the Critical Comments.
- ◆ For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Seedling drench application (brassicas and lettuce)

Ensure even application

Deliver the target rate of Confidor to seedling trays, ensuring uniform distribution of Confidor to each transplant cell. This may be achieved by a single (full rate) or consecutive, multiple (part rate) passes of the application mechanism. Confidor should be applied through dedicated nursery equipment e.g. a hydraulic boom or other suitable equipment which has been carefully set up and calibrated to ensure uniform distribution. **DO NOT use watering cans, hand wands, impact sprinklers and other imprecise means of application.**

Avoid run-through from cells

Applied spray volume must be related to the cell media moisture content and cell capacity, to ensure that there is virtually no loss of applied spray as run-through from cell trays during or following the Confidor application.

Prevent spray drift from treatment area

Spray application should be in an enclosed area, or for example, with an adequately shrouded spray boom, to ensure there is no opportunity for spray drift from the treatment site.

Minimise and capture excess spray

The quantity of excess spray, i.e. applied spray mixture which is not confined in the cell medium, must be minimised by appropriate application equipment set-up. Excess spray includes applied spray which misses the target, runs-off or inadvertently runs-through the cells. Any such excess must be captured and should be filtered and returned to the spray tank or stored for collection by a licensed waste treatment service provider.

Avoid losses from watering after treatment

Avoid watering seedlings after application. If watering is necessary it must be done sparingly. **DO NOT allow run-through from the cells.**

Schedule treatment to minimise time interval to planting

Seedling damage may result from Confidor seedling drench treatment, particularly if transplanting does not occur soon after treatment. It is recommended that transplanting occur within 24 hours of treatment, therefore application should be scheduled accordingly, close to the time of seedling dispatch.

Note on Ornamentals

Confidor 200 SC Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

COMPATIBILITY

Confidor 200 SC is compatible with Antracol® and Baycor®. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Bayer Crop Science, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Bayer Crop Science cannot be responsible for the behaviour of such mixtures.

50 L

Mixing

Prior to pouring, shake container vigorously, then add the required quantity of Confidor 200 SC to water in the spray vat while stirring or with agitators in motion.

Application

Refer to CRITICAL COMMENTS for details of each application method.